

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07J 3/00, 9/00, 17/00, 41/00, A61K</b> <b>31/56, 31/585</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/66611</b> <b>(43) International Publication Date:</b> 9 November 2000 (09.11.00)
<b>(21) International Application Number:</b> PCT/US00/11243 <b>(22) International Filing Date:</b> 27 April 2000 (27.04.00)		<b>(74) Agent:</b> TSAO, Y., Rocky; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).	
<b>(30) Priority Data:</b> 60/131,728 30 April 1999 (30.04.99) US 60/191,864 24 March 2000 (24.03.00) US		<b>(81) Designated States:</b> AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
<b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications</b> US 60/131,728 (CIP) Filed on 30 April 1999 (30.04.99) US 60/191,864 (CIP) Filed on 24 March 2000 (24.03.00)		<b>Published</b> <i>With international search report.</i>	
<b>(71) Applicant (for all designated States except US):</b> ARCH DEVELOPMENT CORPORATION [US/US]; The University of Chicago, Suite 405, 5640 South Ellis, Chicago, IL 60637 (US).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LIAO, Shutsung [US/US]; 5632 South Woodlawn Avenue, Chicago, IL 60637 (US). SONG, Ching [CN/US]; Apartment #3B, 5748 South Drexel Avenue, Chicago, IL 60637 (US).			
<b>(54) Title:</b> STEROID DERIVATIVES			
<b>(57) Abstract</b>  Steroid derivatives of this invention interact with nuclear liver X receptor (LXR) and ubiquitous receptor (UR) and can be used to treat a variety of LXR- or UR-mediated disorders.			

## **STEROID DERIVATIVES**

### **BACKGROUND**

Nuclear receptors are a family of transcription factors modulated by small hydrophobic signaling molecules, like steroids, thyroid hormone, free fatty acids, vitamin D and retinoids. Nuclear receptors are important pharmacological targets for drug intervention in disease management. For example, Tamoxifen, an estrogen antagonist, interacts with estrogen receptor to deliver its therapeutic effects on breast cancer; RU486, an antagonist of progesterone receptor, is used for termination of pregnancies and menopause-related disorders; and Dexamethasone interacts with glucocorticoid receptor to suppress immune system function and is useful for treating inflammatory diseases such as asthma.

Nuclear receptors have three independent domains I, II and III. Domains I and III modulate transcriptional activities by interacting with other factors of the transcription complex; Domain II involves in DNA-binding; and Domain III is the ligand-binding domain. Domain II is the most conserved region within the nuclear receptor family, with a unique feature of four pairs of cysteine chelated with two zinc atoms which form a "zinc finger" structure. The three domains of nuclear receptors are functionally interchangeable between different members. For example, the androgen receptor DNA-binding domain can be fused to the ligand-binding domain of estrogen receptor and the resulting AR-ER chimeric receptor can modulate androgen-responsive genes by binding to estrogen.

Amino acid sequence homology of the DNA-binding domain between members of nuclear receptor family allows identification of new members of this family through low stringency nucleotide-probe screening. Human genome project also facilitates identification of new genes coding for new nuclear receptors. At present, a few dozens of nuclear receptors have been identified and sequenced, but their ligands have yet to be identified. Recently, a novel nuclear

receptor was cloned through degenerate oligonucleotide screening from human and rat cells and was named ubiquitous nuclear receptor ("UR"), because of its ubiquitous expression pattern in the body. UR has been found to form heterodimers with RXR receptors and binds to double-stranded DNA with the sequence motif: AGGTCANNNNAGGTCA (SEQ ID NO: 1) ("DR4"). Promoters containing DR4 can be activated by UR and RXR heterodimer in cultured cells.


LXR $\alpha$ , another new member of the nuclear receptor family has been cloned recently. Amino acid sequence analysis revealed that it shares over 80% homology with UR in the DNA- and ligand-binding domain. The expression of LXR $\alpha$  mRNA is limited to liver and a few other tissues. LXR $\alpha$  has been identified as a transcriptional activator of the cholesterol 7 $\alpha$ -hydroxylase gene and plays an important role in cholesterol catabolism.

Recently other nuclear proteins interacting with nuclear receptors have been identified through yeast two-hybrid screening techniques, among which are co-activators and co-repressors of nuclear receptors, e.g., SRC1, 2, 3, and Grip1. These proteins interact with nuclear receptors in a ligand-dependent manner. This property is useful to set up biochemical assays for ligand-receptor interaction.

Steroid derivatives described in this invention are found to modulate the transcriptional activities via binding to UR or LXR $\alpha$ , and thus can be used to treat disorders mediated by such receptors such as atherosclerosis.

## SUMMARY

An aspect of this invention relates to steroid derivatives of formula (I):



R<sup>3</sup> is hydrogen, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with

5 hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid. Each of R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17'</sup>, independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-

10 CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid. Each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino. R<sup>17</sup> is -X-Y-Z. X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a

15 cyclic moiety with R<sup>16</sup> and the 2 ring carbon atoms to which R<sup>16</sup> and R<sup>17</sup> are bonded. Y is

-CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond. Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B. A being a side chain of an amino acid, and B is hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl. n is 0, 1, or 2. Note that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl.

Another aspect of this invention relates to steroid derivatives having the formula (I) as depicted above. Each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17</sup>, independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid. Each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino. R<sup>17</sup> is -X-Y-Z. X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with R<sup>16</sup> and the 2 ring carbon atoms to which R<sup>16</sup> and R<sup>17</sup> are bonded. Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond. Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or

heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B. A is an amino acid side chain containing an aromatic moiety, and B is hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl. n is 0, 1, or 2. Note that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl.

A further aspect of this invention relates to steroid derivatives of formula (I), *supra*. Each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17</sup>, independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-,  
5 -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-,  
-CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-,  
and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid. Each of R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino. R<sup>17</sup> is -X-  
10 Y-Z. X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with R<sup>16</sup> and the 2 ring carbon atoms to which R<sup>16</sup> and R<sup>17</sup> are bonded. Y is














-CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond. Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B. A is a side chain of an amino acid, and B is hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl. n is 0, 1, or 2. Note that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl; and that at least one of R<sup>3</sup> and R<sup>4</sup>, R<sup>4</sup> and R<sup>5</sup>, R<sup>5</sup> and R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup>, R<sup>12</sup> and R<sup>13</sup>, and R<sup>15</sup> and R<sup>16</sup>, independently, is deleted to form a double bond. One subset of the just-described steroid derivatives encompasses compounds which are featured by the presence of at least one double bond in the rings, which are formed by deleting one or more of the following pairs of substituents: R<sup>3</sup> and R<sup>4</sup>, R<sup>4</sup> and R<sup>5</sup>, R<sup>12</sup> and R<sup>13</sup>, and R<sup>15</sup> and R<sup>16</sup>. Another subset encompasses compounds which are featured by that Z is alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and optionally substituted with hydroxy, alkoxy, amino, or halo; or is -CH(A)-B. A and B are as described above.


Note that X and Z optionally join together to form a cyclic moiety. For example, if both X and Z are alkyl, and Y is -C(=O)-O-, a lactone results from joining X and Z.





A salt of the steroid derivative of this invention is also within the scope of this invention and can be formed, for example, between the steroid of this

invention having a carboxylate and a cationic counterion such as an alkali metal cation, e.g., a sodium ion or a potassium ion; or an ammonium cation that can be substituted with organic groups, e.g., a tetramethylammonium ion or a diisopropyl-ethylammonium ion. A salt of this invention can also form between  
5 the steroid derivative of this invention having a protonated amino group and an anionic counterion, e.g., a sulfate ion, a nitrate ion, a phosphate ion, or an acetate ion.


Set forth below are some examples of steroid derivatives of this invention:







and



As used herein, the term "alkyl" in this disclosure denotes a straight or branched hydrocarbon chain containing 1-8 carbon atoms. Some examples of an alkyl group are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, heptyl, octyl, or 2-methylpentyl. By the term "cycloalkyl" is meant a cyclic hydrocarbon chain that has 3-8 carbon atoms. The cycloalkyl groups herein described may also contain fused rings. Fused rings are rings that share a common carbon-carbon bond. Examples of cycloalkyl groups include, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, and norbornyl.

The term "alkenyl" refers to a straight or branched hydrocarbon chain containing 2-8 carbon atoms and characterized in having one or more double bonds. Examples of a typical alkenyl include, but not limited to, allyl, propenyl,

2-butenyl, 3-hexenyl and 3-octenyl groups. The term "cycloalkenyl" is meant a cyclic hydrocarbon chain containing 3-8 carbon atoms and having at least one or more double bonds. Similar to the definition of cycloalkyl groups above, cycloalkenyl groups may also contain fused rings. Some examples of cycloalkenyl groups are cyclopentenyl, cyclohexenyl, cycloheptenyl, norbornylenyl, and cyclooctenyl groups.

The term "alkynyl" refers to a straight or branched hydrocarbon chain containing 2-8 carbon atoms and characterized in having one or more triple bonds. Some examples of a typical alkynyl are ethynyl, 2-propynyl, and 3-methylbutynyl.

The terms "heterocycloalkyl" and "heterocycloalkenyl" refer to cycloalkyl and cycloalkenyl groups which contain one or more heteroatoms, such as, nitrogen, oxygen, or sulfur. Typical heterocycloalkyl and heterocycloalkenyl groups include tetrahydrofuranyl, tetrahydropyranyl, piperidiny, morpholino, pyrrolinyl and pyrrolidinyl.

"Aryl" represents an aromatic moiety which contains 6-12 carbon atoms and can contain fused rings. A fused ring is an aromatic group which contains at least two aryl rings sharing a common carbon-carbon bond. Typical examples of aryl include phenyl and naphthyl.

"Heteroaryl" groups in this disclosure are aromatic groups containing 5 to 12 ring atoms, in which one or more of these ring atoms are heteroatoms as defined above. Some examples of heteroaryl groups are pyridyl, pyrazinyl, furyl, pyrrolyl, thienyl, thiazolyl, benzimidazolyl, and imidazolyl.

The positions of substituents on each of the cyclic groups described herein may be at any available position, unless specified otherwise. For example, a methyl substituent on a benzene ring can be attached at the *ortho*, *meta*, or *para* position.

The term "alkoxy" is defined as the moiety "-O-alkyl." Some examples are methoxy, ethoxy, propoxy, isopropoxy, and t-butoxy. "Halo" represents a

halogen atom, such as, fluoro, chloro, bromo, or iodo. By the terms "haloalkyl" and "hydroxyalkyl" are meant alkyl groups which are respectively substituted with one or more halogen atoms and one or more hydroxy groups. The nitrogen atom in an amino or amido group present in a steroid derivative of this invention  
5 can be mono- or di-substituted with an alkyl, a cycloalkyl, a heterocycloalkyl, an aryl, or a heteroaryl.

For convenience, a divalent moiety is named herein as if it were a monovalent moiety. For example, "alkyl," such as CH<sub>3</sub>, which is assigned to X, actually stands for "alkylene," such as -CH<sub>2</sub>-. As recognized by a skilled person  
10 in the art, steroid derivatives described herein contain stereocenters. Both the racemic mixtures of isomers and the optically pure isomers are within the scope of this invention.

Yet another aspect of this invention relates to a pharmaceutical composition for treating a UR- or LXRA-mediated disorder which contains a  
15 pharmaceutically acceptable carrier and an effective amount of one or more of the steroid derivatives described above. The use of such a steroid derivative or a salt thereof for the manufacture of a medicament for treating the above-mentioned disorders is also within the scope of this invention.

A still further aspect of this invention relates to a pharmacological  
20 composition for treating cancer, including solid tumors and leukemia, and immune dysfunction. The pharmacological composition contains a pharmaceutically acceptable carrier and an effective amount of one or more of a steroid derivative of formula (I), *supra*. Each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17</sup>, independently, is hydrogen, hydroxy, amino, carboxyl,  
25 oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid. Each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is

hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino.  $R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-,  
5 -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, alkyloxycarbonyl, alkylcarbonyloxy,  
10 alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A being a side chain of an amino acid, and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl; and n is 0, 1, or 2. When Z is substituted with carboxyl, Y is a bond and either X or Z contains at least one double bond,  
15 and when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl. The use of a just-described steroid derivative or a salt thereof for the manufacture of a medicament for treating the above-  
20 mentioned disorders is also within the scope of this invention.

Still another aspect of the present invention relates to a method of treating a UR- or LXRA-mediated disorder by administering to a patient in need thereof an effective amount of one of the pharmaceutical compositions described above. Some examples of UR- or LXRA-mediated disorders are: liver cirrhosis, gallstone  
25 disease, hyperlipoproteinemias, Alzheimer's disease, anemia, chronic inflammatory diseases (e.g., rheumatoid arthritis), metabolic disorders (e.g., diabetes), and cancers which are associated with UR expression, e.g., breast cancer, colon cancer, prostate cancer, and leukemia. Patients with other disorders




such as atherosclerosis and liver cholestasis can also be treated with one of the pharmaceutical compositions described above.

Other features or advantages of the present invention will be apparent from the following detailed description of several embodiments, and also from  
5 the appending claims.

### **DETAILED DESCRIPTION**

A steroid derivative of this invention can be prepared by forming an amide bond between a steroid having a C17 carboxyl-containing substituent and  
10 an amino-containing compound or between a steroid having a C17 amino-containing substituent and a carboxyl-containing compound. Similarly, an ester bond can be formed between a steroid with a C17 carboxyl-containing substituent and a hydroxyl-containing compound, or between a steroid with a C17 hydroxyl-containing substituent and a carboxyl-containing compound. Some examples of a  
15 steroid that can be used as a starting material are cholic acid (e.g., ursodeoxycholic acid, hyocholic acid, and hyodeoxycholic acid), androstan-17-carboxylic acid (e.g., androstan-3-oxo-17-carboxylic acid and d5-androsten-3-ol-17-carboxylic acid) and pregnan-20-ol (e.g., d5-pregnen-3,17-diol or pregnan-17-ol-3-one). Synthesis of these steroids can be found in the literature, e.g., Roda A.  
20 et al., *F. Lipid Res.* vol. 35, pages 2268-2279 (1994) and Roda A. et al., *Dig. Dis. Sci.* vol. 34, pages 24S-35S (1987). Some examples of compounds that can be used to couple to a steroid to form a steroid derivative of this invention are aniline, glycine, phenylalanine, or benzoic acid. Examples of a coupling reagent that can be used in the amide- or ester-forming reaction include 1-ethyl-3-[3-  
25 dimethylaminopropyl]-carbodiimide (EDC), dicyclohexyl-carbodiimide (DCC), N-hydroxybenzo-triazole (HOBt), 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (HBTU), or benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBOP). The amide- or ester-forming reaction can take place in any solvents that are suitable with the

starting materials and reagents. Note that if the reaction takes place in an aqueous solvent, e.g., a buffered solution (or in combination with other miscible organic solvents such as alcohol), isolation of the steroid product for *in vitro* or *in vivo* screening assays is not necessary, as the product is already in suitable assaying conditions, i.e., in an aqueous buffered medium. Protection of functional groups, e.g., hydroxyl or keto, on the steroids is not needed. See, e.g., Example 1 below. Due to the simplicity of the reaction, it can be easily automated. Isolation and quantification of the product can be done by thin-layer chromatography, high pressure liquid chromatography, gas chromatography, capillary electrophoresis, or other analytical and preparative procedures. Trifluoromethyl- and taurine-conjugated steroid derivatives can be prepared according to methods described in Li, S. et al., Chem. Phys. Lipids 99:33-71 (1999) and Kurosawa, T. et al., Steroids, 60:439-444 (1995), respectively. As to the preparation of 3 $\beta$ -hydroxy-5-cholesten-25(R)-26-carboxylic acid derivatives, see Kim, H. et al., J. Lipid Res. 30:247 (1989) and Varma, R. K. et al., J. Org. Chem. 40:3680 (1975). Steroid derivatives having a side chain that contains a double bond, e.g., between C24 and C25, can be prepared according to the following scheme:



3-beta-t-butyldimethylsilyloxy-delta[5]-cholen-24-al and 3-alpha,6-alpha-di(t-butyldimethylsilyloxy)5-beta-cholan-24-al were prepared using NaBH<sub>4</sub> and pyridinium chlorochromate according to methods described in Somanathan et al.,

5 Steroids 43:651-655 (1984). Ethyl-3-beta-t-butyldimethylsilyloxy-delta[5,24]-cholestenoate and ethyl-3a,6a-di(t-butyldimethylsilyloxy)-delta[24]-cholestanoate were then prepared via Wittig-Horner reaction using triethyl 2-phosphono-

propionate and a suitable base according to methods described in Lund et al., Arterioscler. Thromb. Vasc. Biol. 16:208-212 (1996). After the t-

butyldimethylsilyloxyl groups were removed, ethyl ester groups were hydrolyzed under alkaline conditions.

As mentioned above, a pharmaceutical composition containing a steroid derivative or a salt of this invention in an effective amount can be used to treat  
5 UR- or LXRa-mediated disorders. Also within the scope of this invention is a method of treating a UR- or LXRa-mediated disorder such as atherosclerosis by administering to a patient such a composition. An effective amount is defined as the amount of the derivative which, upon administration to a patient in need, confers a therapeutic effect on the treated patient. The effective amount to be  
10 administered to a patient is typically based on body surface area, patient weight, and patient condition. The interrelationship of dosages for patients (based on milligrams per meter squared of body surface) is described by Freireich et al., Cancer Chemother. Rep. 1966, 50, 219. Body surface area may be approximately determined from height and weight of the patient. See, e.g., Scientific Tables,  
15 Geigy Pharmaceuticals, Ardley, New York, 1970, 537. An effective amount of a compound of this invention used to practice the invention can range from about 1 mg/kg to about 2 g/kg, e.g., from about 1 mg/kg to about 1 g/kg, or from about 1 mg/kg to about 500 mg/kg. Effective doses will also vary, as recognized by those skilled in the art, dependant on route of administration, excipient usage, and the  
20 possibility of co-usage with other therapeutic treatments.

The pharmaceutical composition may be administered via the parenteral route, including subcutaneously, intraperitoneally, intramuscularly and intravenously. Examples of parenteral dosage forms include aqueous solutions of the active agent, in a isotonic saline, 5% glucose or other well-known  
25 pharmaceutically acceptable excipient. Solubilizing agents such as cyclodextrins, or other solubilizing agents well-known to those familiar with the art, can be utilized as pharmaceutical excipients for delivery of the therapeutic compounds.

The steroid derivatives of this invention can also be formulated into dosage forms for other routes of administration utilizing well-known methods.

They can be formulated, for example, in dosage forms for oral administration in a gel seal, a syrup, a capsule, or a tablet. Capsules may comprise any well-known pharmaceutically acceptable material such as gelatin or cellulose derivatives. Tablets may be formulated in accordance with the conventional procedure by  
5 compressing mixtures of the compound of this invention and a solid carrier, and a lubricant. Examples of solid carriers include starch and sugar bentonite. The steroid derivatives of this invention can also be administered in a form of a hard shell tablet or a capsule containing a binder (e.g., lactose or mannitol) and a conventional filler.

10           The level of interaction between the UR or LXRA protein and a steroid derivative of this invention can be preliminarily evaluated using various assays as described below:

          Protease protection assay is a simple assay for measuring the level of interaction between a test steroid and the UR or LXRA protein. This assay can be  
15 done by using a <sup>35</sup>S-Met radiolabeled rat UR or human LXRA protein. The radiolabeled protein is then incubated with the steroid of this invention and digested with a protease, e.g., trypsin. A control experiment is done by incubating UR receptor with a protease but without the steroid. Protein fragments from both assays are electrophoresed on a polyacrylamide gel. The  
20 fragments from each of the assays can be visualized by exposing the gel to X-ray films and compared side-by-side. A test steroid, if binds to the UR or LXRA protein, will protect the receptor from being digested by the protease. As a result, reactions that result in binding between the steroid and UR will lead to fewer bands of low molecular weights than those that do not result in binding between  
25 the two molecules.

          The co-activator binding assay employs a fusion protein formed between a glutathione S-transferase (GST) and a co-activator of UR, e.g., Grip1. The GST moiety of the fusion protein binds to a glutathione-coated solid support, thereby immobilizing the fusion protein. UR and a steroid of this invention are

then incubated with the immobilized fusion protein. Subsequently, any bound UR is released and collected from the solid support. The proteins are then electrophoresed on a polyacrylamide gel and visualized by exposing the gel to X-ray films. If the steroid interacts with UR, less UR will bind to the fusion protein, and a lighter band would therefore result on the gel. By monitoring the intensity of the band of the bound UR, one can estimate the binding of the steroid to UR.

Yeast two-hybrid binding assay is a sensitive assay for identifying UR modulating compounds by monitoring transcriptional activation. General descriptions of these assay can be found in, e.g., Chien C. T. et al., *Proc. Natl. Acad. Sci. USA*, vol. 88, 9578-9582 (1991); Fields, S. et al., *Nature*, vol. 340, 245-247 (1989); and Green, M. B. et al., *Curr. Biol.*, vol. 2, 403-405 (1992). In this screening method, a steroid of this invention that modulates the interaction of UR or LXRA with its natural ligand will have an effect on the transcriptional activation of a reporter gene. In a specific assay, two plasmids are introduced into a yeast cell. One expresses a fusion protein having a GAL4 DNA binding domain and a UR natural ligand, and the other expresses a fusion protein containing a UR ligand binding domain and a GAL4 activation domain. If the steroid interacts with UR and disrupts the binding of UR to its natural ligand, the activity of the reporting gene (Gal4) will be altered. The changes in reporter activities (i.e.,  $\beta$ -galactosidase activities) can be measured with a commercial luminescence kit.

Mammalian cell transfection can also be used to screen steroid derivatives that affect the interaction between the UR protein and a steroid of this invention. A rat UR or human LXR gene and a human RXRA gene are cloned into a mammalian expression vector (e.g., pSG5 from Strategene) and overexpressed. A heterologous promoter is formed by inserting four tandem repeats of a hormone response element DR4 into the vector upstream to a c-fos promoter sequence, which is followed by a sequence encoding luciferase. The entire

ARE-fos-luc was constructed by inserting into the vector three tandem repeats of Androgen Response Element (ARE) with a sequence 5'- TCG AGT CTG GTA CAG GGT GTT CTT TTG-3' (SEQ ID NO: 11) upstream to a c-fos promoter sequence (-56-+109) which was followed by a sequence encoding luciferase.

5           Various steroid derivatives of this invention were found to modulate UR transactivation activity on DR4-fos-luc expression in the cultured cells. For example, steroid derivative (6) (see page 6, *supra*) unexpectedly increased the luciferase activity by 5-fold in comparison to the steroid starting material.

10       Mammalian cell transfection assay (3)

Human embryonic kidney 293 cells were seeded into 48-well culture plates at 105 cells per well in DMEM supplemented with 10% fetal bovine serum. After 24 hours, cells were transfected by a calcium phosphate coprecipitation method with 250 ng of the pGL3/UREluc reporter gene which  
15       consists of three copies of AGGTCAagccAGGTCA fused to nucleotides -56 to +109 of the human c-fos promoter in front of the firefly luciferase gene in the plasmid basic pGL3 (Promega), 40 ng pSG5/hRXRa, 40 ng pSG5/rUR or CMX/hLXR, 10 ng pSG5/hGrip1, 0.4 ng CMV/R-luc (transfection normalization reporter, Promega) and 250 ng carrier DNA per well. Alternatively, 500 ng of  
20       the pGL2/7aluc reporter gene which consists of a single copy of nucleotides -101 to -49 of the rat 7 $\alpha$ -hydroxylase gene fused to the SV40 promoter in front of the firefly luciferase gene in the plasmid basic pGL2 (Promega) was used instead of pGL3/UREluc. This reporter does not have response elements for COUP-TFII or HNF4. In some experiments, 500 ng of the human 7 $\alpha$ -hydroxylase gene reporter,  
25       PH/hCYP7A-135, which consists of a single copy of nucleotides -135 to +24 of the human CYP7A gene fused to the firefly luciferase gene in the plasmid basic pGL3 (Promega), was used instead of pGL2/7aluc. After another 12-24 hours, cells were washed with PBS and refed with DMEM supplemented with 4% delipidated fetal bovine serum. Steroid derivatives dissolved in ethanol were

added in duplicate to the medium so that the final concentration of alcohol was 0.2%. After 24-48 hours, cells were harvested and luciferase activity was measured with a commercial kit (Promega Dual luciferase II) on a Monolight luminometer (Beckton Dickenson). Both LXR and UR form heterodimers with RXR for gene transactivation. The ligand for RXR, 9-cis retinoic acid, is known to activate the LXR/RXR heterodimer but addition of 9-cis retinoic acid to transactivation assays did not change the potency of either  $\Delta^5$  or 6 $\alpha$ -hydroxy steroids for activation of LXR or UR (data not shown). The expression of endogenous LXR and UR (and TR which also binds to a DR4 response element) were apparently low since reporter activation was low in the absence of added expression vectors for LXR or UR. Reporter activation was also low when the DR4 response-element was replaced with a glucocorticoid receptor response element. Each experiment was repeated as least twice to demonstrate reproducibility. Relative light units were about  $2 \times 10^7$  for pGL3/UREluc,  $1 \times 10^6$  for pGL2/7aluc,  $5 \times 10^4$  for PH/hCYP7A-135 and  $5 \times 10^5$  for CMV/R-luc. Purity of synthesized steroid derivatives was verified by thin layer chromatography and structures were confirmed using proton and  $C^{13}$  magnetic resonance spectrometry. 3-Oxo-6 $\alpha$ -hydroxy-5 $\beta$ -cholanoic acid methyl ester, 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid methyl ester, and 3 $\alpha$ ,6 $\alpha$ ,7 $\alpha$ -trihydroxy-5 $\beta$ -cholanoic acid methyl ester were found to be as potent as 3 $\beta$ -hydroxy- $\Delta^5$ -cholanoic acid methyl ester as activators for LXR, with ED<sub>50</sub>'s of about 150 nM. Loss of activity was seen when the 6 $\alpha$ -hydroxy group was changed to a 6 $\beta$  configuration. In contrast to activity with LXR, 3 $\beta$ -hydroxy- $\Delta^5$ -cholanoic acid methyl ester (ED<sub>50</sub> of 130 nM) was more active than 3-oxo-6 $\alpha$ -hydroxy-cholanoic acid methyl ester (ED<sub>50</sub> of 550 nM) and 3 $\alpha$ ,6 $\alpha$ -dihydroxy-cholanoic acid methyl ester (ED<sub>50</sub> of 500 nM) for UR activation.

Using the same assay, ED<sub>50</sub>'s of 6 $\alpha$ -hydroxylated steroids with 24-keto side chains include free and conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid and 3 $\alpha$ ,6 $\alpha$ ,7 $\alpha$ -trihydroxy-5 $\beta$ -cholanoic acid were determined. These steroid



derivatives were found to be more selective activators of LXR than UR. 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid activated LXR with an ED<sub>50</sub> of 17 mM for the free acid and 3 mM for its taurine conjugate. Free and taurine-conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acids activated UR with ED<sub>50</sub> of 55 mM and 11 mM, values three to four times higher than those for LXR. Cholanoic acid derivatives containing trifluoromethyl moiety were also found to be selective activators of LXR.

The ability of taurine-conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid to activate LXR using the natural response element derived from the rat 7 $\alpha$ -hydroxylase promoter was also investigated. It was found that taurine-conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid activated LXR but not UR using this reporter gene, with an ED<sub>50</sub> of 10 mM. To investigate if LXR can activate human CYP7A gene transcription, a chimeric reporter plasmid, in which the nucleotides -135 to +24 of the human CYP7A promoter were fused to the luciferase gene, was used in a co-transfection assay in human embryonic kidney 293 cells along with LXR, RXR and Grip1 expression plasmids. It was found that LXR can activate reporter gene expression in the presence of taurine-conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid. Taurine-conjugated 3 $\alpha$ ,7 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid, on the other hand, suppressed reporter gene expression. Another compound, 3 $\beta$ -hydroxy-5-cholesten-25(R)-26-carboxylic acid activated LXR with an ED<sub>50</sub> of 300 nM and UR with an ED<sub>50</sub> of over 2  $\mu$ M. Its taurine-conjugated counterpart was also found to be able to transactivate both LXR and UR. On the other hand, many of its related metabolites were found to be inactive on either receptors.

#### Protease Protection Assay

Rat UR protein radio-labeled with <sup>35</sup>S-Met is produced with a commercial kit in an in vitro system. The radio-labeled protein is incubated with steroid derivatives with final concentration of up to 1 mM for 2 hours on ice, and

digested with trypsin for 30 minutes at 37°C for 20 minutes. The protected fragments were separated from free <sup>35</sup>S-Met by polyacrylamide electrophoresis and visualized by exposing dried gels to X-ray films.

The patterns of the X-ray film indicate that steroid derivatives of this invention bind to and protect UR from being digested by trypsin. Some examples of such a steroid derivative include 5β-androstan-3a,17b-diol, 5β-androstan-3a-ol-16-one, Δ<sup>5</sup>-Pregnen-3b-ol-20-one, 5a-androstan-3-one, 5α-androstan-17-ol-3-one, 5a-androstan-3b-ol-17-carboxylic acid, 5a-pregnan-3,20-dione, and Δ<sup>5</sup>-androst-3b,17b-diol.

Incubation of UR with increasing concentrations of trypsin in the absence of 3α,6α-dihydroxy-5β-cholanoic acid methyl ester leads to extensive digestion of the receptor. In contrast, when UR was incubated with 5 mM 3α,6α-dihydroxy-5β-cholanoic acid methyl ester, two protease-resistant fragments of 35 and 26 kDa were observed. A similar protected pattern was observed with taurine-conjugated 3α,6α-dihydroxy-5β-cholanoic acid.

#### Co-activator Binding Assay

A fusion protein formed between glutathione S-transferase and Grip1 (termed GST-Grip1) was expressed in *E. Coli*. The bacteria was lysed by sonication in the presence of detergent NP40 0.1% and Tween-20 0.5%. Soluble GST-Grip1 was separated from insoluble debris by centrifugation at 50,000G at 4°C for 30 minutes. The soluble fusion protein was then immobilized to glutathione-agarose. Radiolabeled rat UR protein was incubated with GST-Grip1 coated glutathione-agarose in the presence of a test compound of this invention for 2 hours at 22°C under agitation. UR that did not bind to the agarose was washed away. Bound UR was eluted with solution containing SDS and β-mercaptoethanol and separated from free <sup>35</sup>S-Met with polyacrylamide electrophoresis, and finally visualized by exposure the dried gel to X-ray films.

Diosgenin was shown to be capable of promoting UR and Grip 1 protein interaction in this assay.

Another fusion protein GST-rUR was expressed in *E. Coli* strain BL21 using the expression plasmid pGEX using a method similar to that as described above. Transfected cells were lysed by one cycle of freeze-thaw and sonication. Supernatant, prepared by centrifugation at 45,000G for 1 hour, was incubated with glutathione-agarose for 10 min at 4°C. The agarose was washed with binding buffer (20mM Hepes, pH7.5, 10 mM EDTA, 10mM Na<sub>2</sub>MoO<sub>4</sub>, 1 mM β-mercaptoethanol, 1 mM DTT, 0.5 mM PMSF, 2 ug/ml aprotinin). Human Grip1 was produced by in vitro translation using a rabbit reticulocyte lysate and labeled with [<sup>35</sup>S]-methionine. [<sup>35</sup>S]-Grip1 in reticulate lysate (2 ml) was added to GST-UR bound to agarose beads in 100 ul binding buffer. Test chemicals in ethanol were added to the mixture and the slurry was shaken at room temperature for 30 min. The agarose beads were then washed three times with binding buffer. Bound protein was eluted with SDS-PAGE loading buffer and separated on a 8% SDS-PAGE gel. Gels were dried and subjected to autoradiography. Radioactive Grip1 was measured with a STORM phosphoimager (Molecular Dynamics).

Both 3α,6α-dihydroxy-5β-cholanoic acid methyl ester and 22R-hydroxy cholesterol promoted interaction of Grip1 with GST-UR and taurine-conjugated 3α,6α-dihydroxy-5β-cholanoic acid promoted interaction of Grip1 with GST-LXR. Taurine-conjugated 3α-hydroxy-5β-cholanoic acid, taurine-conjugated 3α-hydroxy-5β-cholanoic acid, and taurine-conjugated 3α,7α-dihydroxy-5β-cholanoic acid all failed to enhance coactivator-receptor interaction under the same conditions.

Using the same conditions, 3β-hydroxy-5-cholesten-25(R)-26-carboxylic acid was found to bind to and form complexes with LXR and nuclear receptor co-activator Grip 1, indicating that this acid bound to LXR and induced a conformation change that favored co-activator binding. In a dose response analysis, 3β-hydroxy-5-cholesten-25(R)-26-carboxylic acid increased the amount

of [<sup>35</sup>S]-Grip1 bound to LXR with an EC<sub>50</sub> value of 300 nM, which correlates with the cell-based transfection assay. These data showed that 3β-hydroxy-5-cholesten-25(R)-26-carboxylic acid is a LXR agonist.

5     Inhibition of de novo  
      cholesterol synthesis in cultured cells

On day 1, PC-3 cells stably expressing rUR gene by retroviral infection were seeded in media containing delipidated serum. On day 2, cells were treated with an ethanol solution containing a test compound at maximum concentration of 2  
10    μM. On day 3, cells were washed with PBS and treated with 100 mg/ml amphotericin B in Dulbecco's Modified Eagle Medium (DMEM) without serum at 37°C. 4 hours later, cells were then washed and treated with solution containing 80% water and 20% DMEM for 30 minutes. Surviving cells were assessed using a colorimetric assay. Cells were fixed in 10% trichloroacetic acid  
15    (TCA) and stained with sulforhodamine B. The amount of dye is linear to number of fixed cells on the culture plates. Cells with cholesterol in the cell membrane were killed by amphotericin B treatment.

Compounds of this invention were found to inhibit cholesterol synthesis of the cell to various extent.

20

Measuring the level of inflammation  
      in cells by monitoring the amount of NO<sub>2</sub>

Murine macrophage cell line RAW264.7 were incubated with a test compound at maximum final concentration of 2 μM for 24 hours. The  
25    macrophages were then activated by adding lipopolysaccharide (100 ng/mL) and γ-interferon (100 units/mL). The nitrogen monoxide (NO) production of activated macrophages was measured indirectly by quantifying nitrogen dioxide (NO<sub>2</sub>) in the media according to Green L. et al., *Anal. Biochem.* 126, 131-138 (1982). Compounds of this invention were found to inhibit cholesterol synthesis  
30    of the cell to various extent.

#### Macrophage-foam cell transformation

Constitutive expression of rat UR and human RXRa gene by retroviral systems in RAW264.7 transformed these cells into foam-cell-like morphology and integrated into clamps while increasing cell sizes and undergoing apoptosis. Foam cells originated from macrophages are the major components in pathological plaques formed on the inner wall of blood vessels which are a characteristic feature in atherosclerosis. Compounds of this invention were shown to be able to suppress the progression of macrophage-foam cell transformation at different stages, and thus can be used in the treatment or prevention of atherosclerosis.

#### Adipocyte differentiation

Constitutive expression of rat UR gene in murine fibroblasts 3T3-L1 was done by using retroviral systems. Full-length rat UR cDNA was inserted into retroviral expression vector MV7. Infected 3T3-L1 cells that are G418-resistant were treated with 5 µg/ml insulin, 250 nM dexamethacine, and 0.5 mM 1-methyl-3-isobutylxanthine (MIX) to induce adipocyte differentiation. A control experiment was done by inserting human UR cDNA into MV7 in the antisense orientation. Cells infected with hUR-antisense constructs and parent 3T3-L1 cells were also treated with the same insulin cocktail under same cell density. Cells infected with rUR were shown to accumulate more Red oil O positive lipid drops than parent cells, while cells infected with hUR antisense were shown to have less Red oil O positive lipid drops.

#### Erythrocyte differentiation

Constitutive expression of rat UR gene in murine NN10, IW32.1 or IW201 was done by using retroviral systems. Full-length rat UR cDNA was inserted into retroviral expression vector MV7. Infected cells that were G418-

resistant were cultured upto 5 days to induce erythrocyte differentiation. A control experiment was done by using parent MV7 vector. NN10, IW32.1 or IW201 cells infected with parent MV7 construct were also treated with G418 in parallel under same cell density. More cells infected with rUR were shown to  
5 accumulate hemoglobin protein (stained with benzidine) than parent or control cells. When IW32.1/rUR cells were cultured on fibronectin-coated plates, some cells differentiated into mature enucleated reticulocytes.

#### Animal Studies

10 Male Sprague-Dawley rats that were 50 days old were fed a regular chow diet and tap water *ad libitum* for 1 week during acclimatization, and then randomly divided into groups that were given different dietary treatments. Both control and treatment groups were initially fed *ad libitum* a cholesterol-enriched diet, which was prepared by adding 2% cholesterol and 1% 3 $\alpha$ ,7 $\alpha$ ,12 $\alpha$ -  
15 trihydroxy-5 $\beta$ -cholanoic acid to the regular chow diet. The treatment group received the same diet supplemented with 0.03% test steroid derivative. Rats were fasted overnight before determining body and liver weight and drawing blood from the tail vein for serum total cholesterol measurements. Total cholesterol was determined enzymatically with a diagnostic kit (Sigma, St. Louis,  
20 MO) on day 0 and 7. Average food consumption was 20-25 g/rat/day and average feces production was 9 g/rat/day. There was no statistical difference between control and treatment groups for food consumption and feces production. The dose for test steroid derivative in the treatment group was 40-50  
25 mg/kg/day. Rats on high cholesterol/bile acid diet and treated with a trifluoromethyl conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid had a 20% drop (p < 0.05) in the serum total cholesterol compared with the level in untreated animals (Table 1). Food consumption, body and liver weight were similar in the control and treatment groups. In another experiment, rats were made hypercholesterolemic with a high cholesterol/cholic acid diet and then treatment

with the same trifluoromethyl conjugated 3 $\alpha$ ,6 $\alpha$ -dihydroxy-5 $\beta$ -cholanoic acid again lowered the serum total cholesterol by 20% compared with untreated animals.

5


#### Other Embodiments

From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. For example, the moiety A  
10 can be a side chain of an amino acid which is structurally similar to a naturally occurring amino acid described above. One specific example of A is a side chain of phenylglycine. Thus, other embodiments are also within the claims.

WHAT IS CLAIMED IS:

15

1. A compound of the following formula:



wherein

$R^3$  is hydrogen, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^4'$ ,  $R^6$ ,  $R^7$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{15}$ ,  $R^{16}$ , and  $R^{17}$ , independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of  $R^5$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{13}$ , and  $R^{14}$ , independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;

$R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are



bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A being a side chain of an amino acid, and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl; and

n is 0, 1, or 2;






provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl; or a salt thereof.

2. The compound of claim 1, wherein n is 0.
3. The compound of claim 1, wherein R<sup>3</sup> is amino, carboxyl, halo, sulfonic acid, -O-sulfonic acid, or alkyl; R<sup>6</sup> is hydroxy, amino, carboxyl, halo, sulfonic acid, -O-sulfonic acid, or alkyl; and each of R<sup>3</sup> and R<sup>6</sup>, independently, is in the  $\alpha$ -configuration.
4. The compound of claim 1, wherein R<sup>5</sup> is hydrogen and is in the  $\beta$ -configuration.


5. The compound of claim 1, wherein R<sup>3</sup> is oxo; each of R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17'</sup>, independently, is hydrogen, hydroxy, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl.
6. The compound of claim 5, wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17'</sup>, independently, is hydrogen, hydroxy, or oxo; and each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen or hydroxy; or a salt thereof.
7. The compound of claim 6, wherein X is a bond or alkyl.
8. The compound of claim 7, wherein Y is -C(=O)-NH- or -NH-C(=O)-; and Z is -CH(A)-B with A being a side chain of Tyr or Phe, and B being -NR<sup>a</sup>R<sup>b</sup> or -COOR<sup>c</sup>.
9. The compound of claim 1, wherein X is a bond or alkyl.
10. The compound of claim 9, wherein Y is -C(=O)-NH- or -NH-C(=O)-; and Z is -CH(A)-B with A being a side chain of Tyr or Phe, and B being -NR<sup>a</sup>R<sup>b</sup> or -COOR<sup>c</sup>.
11. The compound of claim 6, wherein Y is -CO-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-NH-, -NH-CO-, or a bond.
12. The compound of claim 11, wherein Z is alkyl, alkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, halo, sulfonic acid, carboxyl, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl; or is -CH(A)-B.

13. The compound of claim 1, wherein Z is alkyl or aryl, each of which being optionally substituted with hydroxy; or is -CH(A)-B with A being an amino acid side chain having an aromatic moiety, and B being -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup>.
14. The compound of claim 1, wherein R<sup>17</sup> contains a straight chain having 6-20  
5 chain atoms.
15. The compound of claim 14, wherein R<sup>17</sup> contains a straight chain having 8-16 chain atoms.
16. The compound of claim 1, wherein X is -CH(CH<sub>3</sub>)-CH<sub>2</sub>-, Y is a bond, and Z is -CH<sub>2</sub>-CH=C(R')(CH<sub>3</sub>) with R' being hydroxy, alkoxy, amino, halo,  
10 sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio.


17. The compound of claim 1, wherein said compound is:



or



18. A compound of the following formula:



wherein

each of  $R^1, R^2, R^3, R^4, R^6, R^7, R^{11}, R^{12}, R^{15}, R^{16}$ , and  $R^{17}$ ,  
independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -  
O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-  
5, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -  
CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally  
substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic  
acid;

each of  $R^5, R^8, R^9, R^{10}, R^{13}$ , and  $R^{14}$ , independently, is hydrogen, alkyl,  
10 haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;

$R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally  
inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a  
cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are  
bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -  
15 O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is  
alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl,  
heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally  
substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid,  
carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl,  
20 alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is  
-CH(A)-B with A being an amino acid side chain containing an aromatic moiety,  
and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>,  
independently, is hydrogen or alkyl; and

n is 0, 1, or 2;

25 provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y  
is a bond and either X or Z contains at least one double bond, and that when Y is  
a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-  
, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo,  
sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl;

or a salt thereof.

19. The compound of claim 18, wherein n is 0.

20. The compound of claim 18, wherein each of R<sup>3</sup> and R<sup>6</sup>, independently, is hydroxy, amino, carboxyl, halo, sulfonic acid, -O-sulfonic acid, or alkyl, and  
5 is in the  $\alpha$ -configuration.

21. The compound of claim 18, wherein R<sup>5</sup> is hydrogen and is in the  $\beta$ -configuration.

22. The compound of claim 18, wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17'</sup>, independently, is hydrogen, hydroxy, oxo, halo,  
10 sulfonic acid, -O-sulfonic acid, or alkyl.

23. The compound of claim 22, wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17'</sup>, independently, is hydrogen, hydroxy, or oxo; and each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen or hydroxy.

24. The compound of claim 23, wherein X is a bond or alkyl.


15

15 25. The compound of claim 24, wherein Y is -C(=O)-NH- or -NH-C(=O)-; and Z is -CH(A)-B with A being a side chain of Tyr or Phe, and B being -NR<sup>a</sup>R<sup>b</sup> or -COOR<sup>c</sup>

26. The compound of claim 18, wherein X is a bond or alkyl.

27. The compound of claim 26, wherein Y is -C(=O)-NH- or -NH-C(=O)-; and Z is -CH(A)-B with A being a side chain of Tyr or Phe, and B being -NR<sup>a</sup>R<sup>b</sup> or -COOR<sup>c</sup>
28. The compound of claim 18, wherein Y is -CO-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-NH-, -NH-CO-, or a bond.
29. The compound of claim 28, wherein Z is alkyl, alkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, halo, sulfonic acid, carboxyl, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl; or is -CH(A)-B.
30. The compound of claim 18, wherein Z is alkyl or aryl, each of which being optionally substituted with hydroxy; or is -CH(A)-B with A being an amino acid side chain having an aromatic moiety, and B being -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup>.
31. The compound of claim 18, wherein R<sup>17</sup> contains a straight chain having 6-20 chain atoms.
32. The compound of claim 31, wherein R<sup>17</sup> contains a straight chain having 8-16 chain atoms.
33. The compound of claim 18, wherein X is -CH(CH<sub>3</sub>)-CH<sub>2</sub>-, Y is a bond, and Z is -CH<sub>2</sub>-CH=C(R')(CH<sub>3</sub>) with R' being hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio.

34. The compound of claim 18, wherein said compound is:




35. A compound of the following formula:




47. The compound of claim 35, wherein X is  $-\text{CH}(\text{CH}_3)-\text{CH}_2-$ , Y is a bond, and Z is  $-\text{CH}_2-\text{CH}=\text{C}(\text{R}')(\text{CH}_3)$  with R' being hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio.

48. The compound of claim 35, wherein said compound is:



49. A compound of the following formula:



wherein

each of  $\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4, \text{R}^4', \text{R}^6, \text{R}^7, \text{R}^{11}, \text{R}^{12}, \text{R}^{15}, \text{R}^{16}$ , and  $\text{R}^{17}$ , independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -

CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of  $R^5$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{13}$ , and  $R^{14}$ , independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;

- 5  $R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -
- 10 CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl,
- 15 alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A being a side chain of an amino acid, and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl; and

n is 0, 1, or 2;

- provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y
- 20 is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl; and


- further provided that at least one of  $R^3$  and  $R^4$ ,  $R^4$  and  $R^5$ ,  $R^5$  and  $R^6$ ,  $R^7$
- 25 and  $R^8$ ,  $R^{12}$  and  $R^{13}$ , and  $R^{15}$  and  $R^{16}$ , independently, is deleted to form a double bond;

or a salt thereof.


50. The compound of claim 49, wherein n is 0.

51. The compound of claim 49, wherein  $R^3$  is hydroxy, amino, carboxyl, halo, sulfonic acid, -O-sulfonic acid, or alkyl, and is in the  $\alpha$ -configuration.
52. The compound of claim 49, wherein each of  $R^1, R^2, R^3, R^4, R^4', R^6, R^7, R^{11}, R^{12}, R^{15}, R^{16}$ , and  $R^{17'}$ , independently, is hydrogen, hydroxy, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl.
53. The compound of claim 52, wherein each of  $R^1, R^2, R^3, R^4, R^4', R^6, R^7, R^{11}, R^{12}, R^{15}, R^{16}$ , and  $R^{17'}$ , independently, is hydrogen, hydroxy, or oxo; and each of  $R^5, R^8, R^9, R^{10}, R^{13}$ , and  $R^{14}$ , independently, is hydrogen or hydroxy.
54. The compound of claim 53, wherein X is a bond or alkyl.
55. The compound of claim 54, wherein Y is  $-C(=O)-NH-$  or  $-NH-C(=O)-$ ; and Z is  $-CH(A)-B$  with A being a side chain of Tyr or Phe, and B being  $-NR^aR^b$  or  $-COOR^c$ .
56. The compound of claim 49, wherein X is a bond or alkyl.
57. The compound of claim 56, wherein Y is  $-C(=O)-NH-$  or  $-NH-C(=O)-$ ; and Z is  $-CH(A)-B$  with A being a side chain of Tyr or Phe, and B being  $-NR^aR^b$  or  $-COOR^c$ .
58. The compound of claim 49, wherein Y is  $-CO-, -O-SO_2-, -SO_2-O-, -O-SO_3-, -SO_3-O-, -CO-NH-, -NH-CO-,$  or a bond.
59. The compound of claim 49, wherein  $R^{17}$  contains a straight chain having 6-20 chain atoms.

60. The compound of claim 59, wherein  $R^{17}$  contains a straight chain having 8-16 chain atoms.
61. The compound of claim 49, wherein X is  $-\text{CH}(\text{CH}_3)-\text{CH}_2-$ , Y is a bond, and Z is  $-\text{CH}_2-\text{CH}=\text{C}(\text{R}')(\text{CH}_3)$  with R' being hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio.
62. The compound of claim 49, wherein said compound is:



63. A pharmaceutical composition for treating a UR- or a LXR-mediated disorder, said composition comprising a pharmaceutically acceptable carrier and an effective amount of a compound of the following formula:



wherein

R<sup>3</sup> is hydrogen, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17</sup>, independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of  $R^5$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{13}$ , and  $R^{14}$ , independently, is hydrogen, alkyl, haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;







$R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a  
 5 cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally  
 10 substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A being a side chain of an amino acid, and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or  
 15 alkyl; and

n is 0, 1, or 2;

provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-,  
 20 , -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl; or a salt thereof.

64. The composition of claim 63, wherein said compound

is:




or

65. A pharmaceutical composition for treating a UR- or a LXR-mediated

5 disorder, said composition comprising a pharmaceutically acceptable carrier  
and an effective amount of a compound of the following formula:





wherein

each of  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^4'$ ,  $R^6$ ,  $R^7$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{15}$ ,  $R^{16}$ , and  $R^{17}$ ,  
independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -  
O-sulfonic acid, or alkyl that is optionally inserted with -NH-, -N(alkyl)-, -O-, -S-,  
5 -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -  
CO-NH-, -CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally  
substituted with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic  
acid;

10 each of  $R^5$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{13}$ , and  $R^{14}$ , independently, is hydrogen, alkyl,  
haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;


$R^{17}$  is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally  
inserted with -NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a  
cyclic moiety with  $R^{16}$  and the 2 ring carbon atoms to which  $R^{16}$  and  $R^{17}$  are  
bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -  
15 O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is  
alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl,  
heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is substituted  
with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo,  
20 alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino,  
alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A

being an amino acid side chain containing an aromatic moiety, and B being hydrogen,  $-NR^aR^b$ , or  $-COOR^c$  wherein each of  $R^a$ ,  $R^b$ , and  $R^c$ , independently, is hydrogen or alkyl; and


n is 0, 1, or 2;

- 5           provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl;
- 10       or a salt thereof.

66. The composition of claim 65, wherein said compound is:



67. A pharmaceutical composition for treating a UR- or a LXR-mediated disorder, said composition comprising a pharmaceutically acceptable carrier and an effective amount of a compound of the following formula:



5 wherein

each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>4'</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>16</sup>, and R<sup>17</sup>,  
independently, is hydrogen, hydroxy, amino, carboxyl, oxo, halo, sulfonic acid, -  
O-sulfonic acid, or alkyl optionally inserted with -NH-, -N(alkyl)-, -O-, -S-, -SO-,  
-SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-, -CO-O-, -O-CO-, -CO-NH-, -  
10 CO-N(alkyl)-, -NH-CO-, or -N(alkyl)-CO-, and further optionally substituted  
with hydroxy, halo, amino, carboxyl, sulfonic acid, or -O-sulfonic acid;

each of R<sup>5</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>13</sup>, and R<sup>14</sup>, independently, is hydrogen, alkyl,  
haloalkyl, hydroxyalkyl, alkoxy, hydroxy, or amino;


R<sup>17</sup> is -X-Y-Z, in which X is a bond, or alkyl or alkenyl, optionally  
15 inserted with

-NH-, -N(alkyl)-, -O-, or -S-, and further optionally forming a cyclic moiety with R<sup>16</sup> and the 2 ring carbon atoms to which R<sup>16</sup> and R<sup>17</sup> are bonded; Y is -CO-, -SO-, -SO<sub>2</sub>-, -O-SO<sub>2</sub>-, -SO<sub>2</sub>-O-, -O-SO<sub>3</sub>-, -SO<sub>3</sub>-O-, -CO-O-, -O-CO-, -CO-NH-, -CO-N(alkyl)-, -NH-CO-, -N(alkyl)-CO-, or a bond; and Z is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, heterocycloalkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl, and is optionally substituted with hydroxy, alkoxy, amino, halo, sulfonic acid, -O-sulfonic acid, carboxyl, oxo, alkyloxycarbonyl, alkylcarbonyloxy, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, or alkylthio; or is -CH(A)-B with A being a side chain of an amino acid, and B being hydrogen, -NR<sup>a</sup>R<sup>b</sup>, or -COOR<sup>c</sup> wherein each of R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup>, independently, is hydrogen or alkyl; and

n is 0, 1, or 2;

provided that when Z is substituted with carboxyl or alkyloxycarbonyl, Y is a bond and either X or Z contains at least one double bond, and that when Y is a bond, either X is -NH-alkyl-, -NH-alkenyl-, -N(alkyl)-alkyl-, -N(alkyl)-alkenyl-, -O-alkyl-, -O-alkenyl-, -S-alkyl-, or -S-alkenyl-; or Z is substituted with halo, sulfonic acid, -O-sulfonic acid, alkylsulfinyl, or alkylsulfonyl, or is alkenyl; and further provided that at least one of R<sup>3</sup> and R<sup>4</sup>, R<sup>4</sup> and R<sup>5</sup>, R<sup>5</sup> and R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup>, R<sup>12</sup> and R<sup>13</sup>, and R<sup>15</sup> and R<sup>16</sup>, independently, is deleted to form a double bond; or a salt thereof.

68. The composition of claim 67, wherein said compound is:



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/11243

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) :C07J 3/00, 9/00, 17/00, 41/00; A61K 31/56, 31/585 US CL :Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 552/522, 542, 546, 547, 553, 554, 555, 610, 611, 641; 540/102; 514/169, 175, 177, 178, 182 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) STN EXPRESS														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X -- Y	XIA, P. et al. Synthesis of N-substituted 3-oxo-17.beta.-carboxamide-4-aza-5.alpha.-androstanes and the Tautomerism of 3-oxo-4-aza-5-androstenes. Heterocycles. April 1998, Vol. 47, No. 2, pages 703-716, especially page 705, compound 10a, see entire document.	35,36,38,42-44 and 67 ----- 39,40												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A* document defining the general state of the art which is not considered to be of particular relevance</td> <td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*E* earlier document published on or after the international filing date</td> <td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>*Δ* document member of the same patent family</td> </tr> <tr> <td>*O* document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>*P* document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Δ* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Δ* document member of the same patent family													
*O* document referring to an oral disclosure, use, exhibition or other means														
*P* document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search		Date of mailing of the international search report												
07 JULY 2000		08 AUG 2000												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>Barbara Badio</i> BARBARA BADIO Telephone No. (703) 308-1235												

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/11243

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	RUELLE et al. The Mobile Order Solubility Equation Applied to Polyfunctional Molecules: The Non-Hydroxysteroids in Aqueous and Non-Aqueous Solvents. International Journal of Pharmaceutics. July 1997, Vol. 157, No. 2, pages 219-232, especially page 222, Table 1 and pages 224-228, Tables 4 and 5, see entire document.	1,2,5,9,13,18,19,22, 26,30,35, 36,38, 42-46,63, 65 and 67 ----- 4,6,7,17,21,23,24, 39, 40 and 64
X -- Y	ANGELICO et al. Dissolution of Human Cholesterol Gallstones in Bile Salt/Lecithin Mixtures: Effect of Bile Salt Hydrophobicity and Various pHs. Scandinavian Journal of Gastroenterology. 1995, Vol. 30, No. 12, pages 1178-1185, see entire document.	18-22,26, 28,29,63, 65 and 67 ----- 23-25